

April 4, 2024



Rigel Announces Publication of Data on REZLIDHIA® (Olutasidenib) in Post-Venetoclax Patients with Mutant IDH1 AML in Leukemia & Lymphoma

- *Olutasidenib induced durable composite complete remission in 43.8% of patients relapsed or refractory to prior venetoclax-based regimens*
- *Safety was consistent with the overall profile of olutasidenib*
- *Olutasidenib may offer a valuable treatment option for patients with mIDH1 previously treated with venetoclax*

SOUTH SAN FRANCISCO, Calif., April 4, 2024 /PRNewswire/ -- Rigel Pharmaceuticals, Inc. (Nasdaq: RIGL) today announced a peer-reviewed publication in *Leukemia & Lymphoma* on data from an analysis of the Phase 2 study evaluating REZLIDHIA® (olutasidenib), a potent, selective, oral, small-molecule inhibitor of mutant isocitrate dehydrogenase-1 (mIDH1)¹, in patients with mIDH1 acute myeloid leukemia (AML) who were relapsed/refractory (R/R) to prior venetoclax-based regimens.

"Venetoclax in combination with a hypomethylating agent is currently standard treatment for patients with newly diagnosed AML who are unfit for intensive chemotherapy, including those with mIDH1. When this therapy fails, patients historically have had limited treatment options and poor prognoses," said Jorge E. Cortes, M.D., Director, Georgia Cancer Center, Cecil F. Whitaker Jr., GRA Eminent Scholar Chair in Cancer, and Phase 2 trial investigator. "The findings from these analyses suggest that REZLIDHIA may provide an effective treatment for patients with AML following failure of venetoclax combination therapy. REZLIDHIA induced durable remissions consistent with those observed in the pivotal trial and had a favorable tolerability profile in this challenging to treat patient population, representing a valuable treatment option."

"These data support REZLIDHIA's efficacy and well-characterized safety profile in patients with mIDH1 R/R AML who had previously been treated with venetoclax combination regimens," said Raul Rodriguez, Rigel's president and CEO. "These analyses are important because they provide valuable insights into the potential benefit of REZLIDHIA in different segments of the mIDH1 R/R AML patient population."

Key points from the paper are summarized below:

- Olutasidenib alone or in combination with azacitidine demonstrated potential efficacy in patients with AML following failure of venetoclax combination therapy
- Of the 18 patients with prior venetoclax treatment, 10 were relapsed, 6 were refractory, and 2 had complete remission with incomplete hematologic recovery (CRi) to a venetoclax combination
- Of the 16 R/R patients, 7 (43.8%) achieved a composite complete remission (CRc), 4 (25%) achieved complete remission (CR), and 1 (6.3%) achieved CR with partial hematologic recovery (CRh). Both patients with CRi at study entry achieved CR
- Median time to CRc was 1.9 months (range 1-2.8). As of the data cut-off (June 18, 2021), median duration of CRc was not reached (range, 1.2-NR, ongoing at 30.4+ months)
- Red blood cell and platelet transfusion independence was achieved in 2/12 (17%) and 2/7 (29%) transfusion-dependent R/R patients at baseline, respectively
- Safety was consistent with the overall profile of olutasidenib

The paper, titled "Olutasidenib in post-venetoclax patients with mutant isocitrate dehydrogenase 1 (*mIDH1*) acute myeloid leukemia (AML)," was published online in *Leukemia & Lymphoma* and can be accessed [here](#).

About AML

Acute myeloid leukemia (AML) is a rapidly progressing cancer of the blood and bone marrow that affects myeloid cells, which normally develop into various types of mature blood cells. AML occurs primarily in adults and accounts for about 1 percent of all adult cancers. The American Cancer Society estimates that there will be about 20,800 new cases in the United States, most in adults, in 2024.²

Relapsed AML affects about half of all patients who, following treatment and remission, experience a return of leukemia cells in the bone marrow.³ Refractory AML, which affects between 10 and 40 percent of newly diagnosed patients, occurs when a patient fails to achieve remission even after intensive treatment.⁴ Quality of life declines for patients with each successive line of treatment for AML, and well-tolerated treatments in relapsed or refractory disease remain an unmet need.

About REZLIDHIA®

INDICATION

REZLIDHIA is indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test.

IMPORTANT SAFETY INFORMATION

WARNING: DIFFERENTIATION SYNDROME

Differentiation syndrome, which can be fatal, can occur with REZLIDHIA treatment. Symptoms may include dyspnea, pulmonary infiltrates/pleuropericardial effusion, kidney injury, hypotension, fever, and weight gain. If differentiation syndrome is suspected, withhold REZLIDHIA and initiate treatment with corticosteroids and hemodynamic monitoring until symptom resolution.

WARNINGS AND PRECAUTIONS

Differentiation Syndrome

REZLIDHIA can cause differentiation syndrome. In the clinical trial of REZLIDHIA in patients with relapsed or refractory AML, differentiation syndrome occurred in 16% of patients, with grade 3 or 4 differentiation syndrome occurring in 8% of patients treated, and fatalities in 1% of patients. Differentiation syndrome is associated with rapid proliferation and differentiation of myeloid cells and may be life-threatening or fatal. Symptoms of differentiation syndrome in patients treated with REZLIDHIA included leukocytosis, dyspnea, pulmonary infiltrates/pleuropericardial effusion, kidney injury, fever, edema, pyrexia, and weight gain. Of the 25 patients who experienced differentiation syndrome, 19 (76%) recovered after treatment or after dose interruption of REZLIDHIA. Differentiation syndrome occurred as early as 1 day and up to 18 months after REZLIDHIA initiation and has been observed with or without concomitant leukocytosis.

If differentiation syndrome is suspected, temporarily withhold REZLIDHIA and initiate systemic corticosteroids (e.g., dexamethasone 10 mg IV every 12 hours) for a minimum of 3 days and until resolution of signs and symptoms. If concomitant leukocytosis is observed, initiate treatment with hydroxyurea, as clinically indicated. Taper corticosteroids and hydroxyurea after resolution of symptoms. Differentiation syndrome may recur with premature discontinuation of corticosteroids and/or hydroxyurea treatment. Institute supportive measures and hemodynamic monitoring until improvement; withhold dose of REZLIDHIA and consider dose reduction based on recurrence.

Hepatotoxicity

REZLIDHIA can cause hepatotoxicity, presenting as increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), increased blood alkaline phosphatase, and/or elevated bilirubin. Of 153 patients with relapsed or refractory AML who received REZLIDHIA, hepatotoxicity occurred in 23% of patients; 13% experienced grade 3 or 4 hepatotoxicity. One patient treated with REZLIDHIA in combination with azacitidine in the clinical trial, a combination for which REZLIDHIA is not indicated, died from complications of drug-induced liver injury. The median time to onset of hepatotoxicity in patients with relapsed or refractory AML treated with REZLIDHIA was 1.2 months (range: 1 day to 17.5 months) after REZLIDHIA initiation, and the median time to resolution was 12 days (range: 1 day to 17 months). The most common hepatotoxicities were elevations of ALT, AST, blood alkaline phosphatase, and blood bilirubin. Monitor patients frequently for clinical symptoms of hepatic dysfunction such as fatigue, anorexia, right upper abdominal discomfort, dark urine, or jaundice. Obtain baseline liver function tests prior to initiation of REZLIDHIA, at least once weekly for the first two months, once every other week for the third month, once in the fourth month, and once every other month for the duration of therapy. If hepatic dysfunction occurs, withhold, reduce, or permanently discontinue REZLIDHIA based on recurrence/severity.

ADVERSE REACTIONS

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were aspartate aminotransferase increased, alanine aminotransferase increased, potassium decreased, sodium decreased, alkaline phosphatase increased, nausea, creatinine increased, fatigue/malaise, arthralgia, constipation, lymphocytes increased, bilirubin increased, leukocytosis, uric acid increased, dyspnea, pyrexia, rash, lipase increased, mucositis, diarrhea and transaminitis.

DRUG INTERACTIONS

- Avoid concomitant use of REZLIDHIA with strong or moderate CYP3A inducers.
- Avoid concomitant use of REZLIDHIA with sensitive CYP3A substrates unless otherwise instructed in the substrates prescribing information. If concomitant use is unavoidable, monitor patients for loss of therapeutic effect of these drugs.

LACTATION

Advise women not to breastfeed during treatment with REZLIDHIA and for 2 weeks after the last dose.

GERIATRIC USE

No overall differences in effectiveness were observed between patients 65 years and older and younger patients. Compared to patients younger than 65 years of age, an increase in incidence of hepatotoxicity and hypertension was observed in patients ≥ 65 years of age.

HEPATIC IMPAIRMENT

In patients with mild or moderate hepatic impairment, closely monitor for increased probability of differentiation syndrome.

[Click here](#) for Full Prescribing Information, including Boxed WARNING.

To report side effects of prescription drugs to the FDA, visit www.fda.gov/medwatch or call 1-800-FDA-1088 (800-332-1088).

REZLIDHIA is a registered trademark of Rigel Pharmaceuticals, Inc.

About Rigel

Rigel Pharmaceuticals, Inc. (Nasdaq: RIGL) is a biotechnology company dedicated to discovering, developing and providing novel therapies that significantly improve the lives of patients with hematologic disorders and cancer. Founded in 1996, Rigel is based in South San Francisco, California. For more information on Rigel, the Company's marketed products and pipeline of potential products, visit www.rigel.com.

1. de Botton S, et al. Olutasidenib (FT-2102) induces durable complete remissions in patients with relapsed or refractory IDH1-mutated AML. *Blood Advances*. February 1, 2023.
doi: <https://doi.org/10.1182/bloodadvances.2022009411>
The American Cancer Society. Key Statistics for Acute Myeloid Leukemia (AML). Revised January 17, 2024. Accessed Feb. 19, 2024:
<https://www.cancer.org/cancer/acute-myeloid-leukemia/about/key-statistics.html>
2. Leukaemia Care. Relapse in Acute Myeloid Leukaemia (AML). Version 3. Reviewed October 2021. Accessed Feb 19, 2024: <https://media.leukaemiacare.org.uk/wp-content/uploads/Relapse-in-Acute-Myeloid-Leukaemia-AML-Web-Version.pdf>
3. Thol F, Schlenk RF, Heuser M, Ganser A. *How I treat refractory and early relapsed acute myeloid leukemia*. *Blood* (2015) 126 (3): 319-27.
doi: <https://doi.org/10.1182/blood-2014-10-551911>

Forward-Looking Statements

This press release contains forward-looking statements relating to, among other things, that

olutasidenib may provide a meaningful approach to the treatment of Post-Venetoclax Patients with Mutant IDH1 AML. Any statements contained in this press release that are not statements of historical fact may be deemed to be forward-looking statements. Forward-looking statements can be identified by words such as "may", "potential", "look forward", "believe", "will" and similar expressions in reference to future periods. Forward-looking statements are neither historical facts nor assurances of future performance. Instead, they are based on Rigel's current beliefs, expectations, and assumptions and hence they inherently involve significant risks, uncertainties and changes in circumstances that are difficult to predict and many of which are outside of our control. Therefore, you should not rely on any of these forward-looking statements. Actual results and the timing of events could differ materially from those anticipated in such forward looking statements as a result of these risks and uncertainties, which include, without limitation, risks and uncertainties associated with the FDA, European Medicines Agency, PMDA or other regulatory authorities may make adverse decisions regarding olutasidenib; risks that clinical trials may not be predictive of real-world results or of results in subsequent clinical trials; risks that olutasidenib may have unintended side effects, adverse reactions or incidents of misuses; the availability of resources to develop Rigel's product candidates; market competition; as well as other risks detailed from time to time in Rigel's reports filed with the Securities and Exchange Commission, including its Annual Report on Form 10-K for the year ended December 31, 2023 and subsequent filings. Any forward-looking statement made by us in this press release is based only on information currently available to us and speaks only as of the date on which it is made. Rigel does not undertake any obligation to update forward-looking statements, whether written or oral, that may be made from time to time, whether as a result of new information, future developments or otherwise, and expressly disclaims any obligation or undertaking to release publicly any updates or revisions to any forward-looking statements contained herein, except as required by law.

Contact for Investors & Media:

Investors:

Rigel Pharmaceuticals, Inc.
650.624.1232
ir@rigel.com

Media:

David Rosen
Argot Partners
212.600.1902
david.rosen@argotpartners.com



View original content to download multimedia <https://www.prnewswire.com/news-releases/rigel-announces-publication-of-data-on-rezlidhia-olutasidenib-in-post-venetoclax-patients-with-mutant-idh1-aml-in-leukemia--lymphoma-302107833.html>

SOURCE Rigel Pharmaceuticals, Inc.