

March 9, 2026



Tonix Pharmaceuticals Presented Data on TONMYA™ in Treating Patients with Fibromyalgia at the 2026 AAPM PainConnect Annual Meeting

TONMYA (cyclobenzaprine HCl sublingual tablets) for the treatment of fibromyalgia in adults, was commercially launched in November 2025

Treatment with TONMYA provided statistically significant pain reduction in two Phase 3 trials and was generally well tolerated

BERKELEY HEIGHTS, N.J., March 09, 2026 (GLOBE NEWSWIRE) -- Tonix Pharmaceuticals Holding Corp. (Nasdaq: TNXP) (“Tonix” or the “Company”), a fully integrated, commercial biotechnology company, presented data on TONMYA™ (cyclobenzaprine HCl sublingual tablets), at the 2026 American Academy of Pain Medicine (AAPM) PainConnect Annual Meeting, in Salt Lake City, Utah.

“The data presented at the AAPM PainConnect Annual Meeting support TONMYA’s role as a safe and effective non-opioid analgesic for daily use at bedtime in fibromyalgia,^{1,2}” said Gregory Sullivan, M.D., Chief Medical Officer of Tonix Pharmaceuticals. “TONMYA is designed to target nonrestorative sleep and is a first-in-class tertiary amine tricyclic for long-term use. As the first approved therapy for fibromyalgia in over 15 years, TONMYA is an alternative to three currently FDA-approved medicines, which are limited by side effects, adherence, and high discontinuation rates. TONMYA’s unique sublingual formulation enables cyclobenzaprine to bypass first-pass hepatic metabolism. Relative to off-label oral swallowed cyclobenzaprine, TONMYA treatment results in reduced formation of the active, persistent metabolite norcyclobenzaprine, which we believe interferes with the durability of cyclobenzaprine’s treatment effect in fibromyalgia with long term dosing. Oral cyclobenzaprine failed in development because it provided only short-term (one month) benefit³, which is not sufficient for the treatment of fibromyalgia, a chronic condition that requires a sustained treatment effect.”

Data presented at the AAPM PainConnect Annual Meeting are from the Phase 3 RESILIENT trial, a 14-week randomized, double-blind, placebo-controlled study that assessed the safety and efficacy of TONMYA in 457 patients who met the 2016 American College of Rheumatology (ACR) diagnostic criteria for fibromyalgia. In addition to showing a statistically significant reduction in mean daily pain at 14 weeks, TONMYA treatment resulted in an increase over placebo in the number of individuals with a 30% reduction in daily pain, which is considered a clinically meaningful response. The most common adverse events were mild and self-limited oral cavity reactions that uncommonly led to study withdrawal.

TONMYA was approved by the FDA in August 2025 in part based on the Phase 3 RESILIENT trial results and was commercially launched in the U.S. in November 2025.

A copy of the Company's poster presentation, "Treatment with TNX-102 SL Produces Clinically Meaningful Improvements in Patient-Centered Outcomes in Fibromyalgia," is available under the Presentations tab of the Tonix website at <https://ir.tonixpharma.com/presentations>.

Citations

¹Carette S, et al. *Arthritis Rheum*. 1994. 37(1):32-40. doi: 10.1002/art.1780370106.

²Lederman S, et al. *Arthritis Care Res (Hoboken)*. 2023. 75(11):2359-2368. doi: 10.1002/acr.25142.

³Lederman S, et al. *Pain Med*. 2026. 27(1):86-94. doi: 10.1093/pm/pnaf089.

About Fibromyalgia

Fibromyalgia is a chronic pain disorder that is understood to result from amplified sensory and pain signaling within the central nervous system. Fibromyalgia afflicts an estimated 6-12 million adults in the U.S., approximately 90% of whom are women. Symptoms of fibromyalgia include chronic widespread pain, nonrestorative sleep, fatigue, and morning stiffness. Other associated symptoms include cognitive dysfunction and mood disturbances, including anxiety and depression. Individuals suffering from fibromyalgia struggle with their daily activities, have impaired quality of life, and frequently are disabled. Physicians and patients report common dissatisfaction with currently marketed products.

About TONMYA™ (cyclobenzaprine HCl sublingual tablets)

TONMYA (cyclobenzaprine HCl sublingual tablets) is a patented sublingual tablet formulation of cyclobenzaprine hydrochloride which provides rapid transmucosal absorption and reduced production of a long half-life active metabolite, norcyclobenzaprine, due to bypass of first-pass hepatic metabolism. As a multifunctional agent with potent binding and antagonist activities at the 5-HT_{2A} serotonergic, α₁-adrenergic, H₁-histaminergic, and M₁-muscarinic receptors, TONMYA was approved on August 15, 2025, by the FDA for the treatment of fibromyalgia in adults. TONMYA is the first new prescription medicine approved for fibromyalgia in more than 15 years. TONMYA was investigated as TNX-102 SL. TNX-102 SL is also being developed to treat acute stress reaction (ASR)/acute stress disorder (ASD), and major depressive disorder (MDD). The United States Patent and Trademark Office (USPTO) issued United States Patent No. 9636408 in May 2017, Patent No. 9956188 in May 2018, Patent No. 10117936 in November 2018, Patent No. 10,357,465 in July 2019, and Patent No. 10736859 in August 2020. The Protectic™ protective eutectic and Angstro-Technology™ formulation claimed in the patent are important elements of Tonix's proprietary TONMYA composition. These patents are expected to provide TONMYA with U.S. market exclusivity until 2034/2035.

Tonix Pharmaceuticals Holding Corp.

Tonix Pharmaceuticals* is a fully-integrated, commercial-stage biotechnology company focused on central nervous system (CNS) and immunology treatments in areas of high

unmet medical need. TONMYA™ (cyclobenzaprine HCl sublingual tablets 2.8 mg), the Company's recently approved flagship medicine, is the first new treatment for fibromyalgia in more than 15 years. Tonix's CNS commercial infrastructure supports its marketed products, including its acute migraine products, Zembrace® SymTouch® and Tosymra®. Tonix is maximizing the science behind TONMYA in Phase 2 clinical trials to evaluate its potential in major depressive disorder and acute stress disorder. In addition, the company's CNS portfolio includes TNX-2900, which is Phase 2 ready for the treatment of Prader-Willi syndrome, a rare disease. Tonix is also advancing a pipeline of immunology programs, including monoclonal antibody TNX-4800 for Lyme disease prophylaxis and TNX-1500, a third-generation CD40 ligand inhibitor for the prevention of kidney transplant rejection. To learn more, visit www.tonixpharma.com and follow the Company on [LinkedIn](#) and [X](#).

*Tonix's product development candidates are investigational new drugs or biologics; their efficacy and safety have not been established and have not been approved for any indication.

Tonmya, Zembrace SymTouch and Tosymra are registered trademarks of Tonix Medicines.

Forward Looking Statements

Certain statements in this press release are forward-looking within the meaning of the Private Securities Litigation Reform Act of 1995 including those relating to the completion of the offering, the satisfaction of customary closing conditions, the intended use of proceeds from the offering and other statements that are predictive in nature. These statements may be identified by the use of forward-looking words such as "anticipate," "believe," "forecast," "estimate," "expect," and "intend," among others. These forward-looking statements are based on Tonix's current expectations and actual results could differ materially as a result of a number of factors, including the ability of the Company to satisfy the conditions to the closing of the offering and the timing thereof, as well as those described in the Company's Annual Report on Form 10-K for the year ended December 31, 2024, as filed with the SEC on March 18, 2025, and periodic reports filed with the SEC on or after the date thereof. Tonix does not undertake an obligation to update or revise any forward-looking statement. All of Tonix's forward-looking statements are expressly qualified by all such risk factors and other cautionary statements. The information set forth herein speaks only as of the date thereof.

Investor Contacts

Jessica Morris
Tonix Pharmaceuticals
investor.relations@tonixpharma.com
(862) 799-8599

Brian Korb
astr partners
(917) 653-5122
brian.korb@astrpartners.com

Media Contacts

Deborah Elson
Tonix Pharmaceuticals

deborah.elson@tonixpharma.com

Ray Jordan
Putnam Insights
ray@putnaminsights.com

INDICATION

TONMYA is indicated for the treatment of fibromyalgia in adults.

CONTRAINDICATIONS

TONMYA is contraindicated: In patients with hypersensitivity to cyclobenzaprine or any inactive ingredient in TONMYA. Hypersensitivity reactions may manifest as an anaphylactic reaction, urticaria, facial and/or tongue swelling, or pruritus. Discontinue TONMYA if a hypersensitivity reaction is suspected. With concomitant use of monoamine oxidase (MAO) inhibitors or within 14 days after discontinuation of an MAO inhibitor. Hyperpyretic crisis seizures and deaths have occurred in patients who received cyclobenzaprine (or structurally similar tricyclic antidepressants) concomitantly with MAO inhibitors drugs. During the acute recovery phase of myocardial infarction, and in patients with arrhythmias, heart block or conduction disturbances, or congestive heart failure. In patients with hyperthyroidism.

WARNINGS AND PRECAUTIONS

Embryofetal toxicity: Based on animal data, TONMYA may cause neural tube defects when used two weeks prior to conception and during the first trimester of pregnancy. Advise females of reproductive potential of the potential risk and to use effective contraception during treatment and for two weeks after the final dose. Perform a pregnancy test prior to initiation of treatment with TONMYA to exclude use of TONMYA during the first trimester of pregnancy.

Serotonin syndrome: Concomitant use of TONMYA with selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, tramadol, bupropion, meperidine, verapamil, or MAO inhibitors increases the risk of serotonin syndrome, a potentially life-threatening condition. Serotonin syndrome symptoms may include mental status changes, autonomic instability, neuromuscular abnormalities, and/or gastrointestinal symptoms. Treatment with TONMYA and any concomitant serotonergic agent should be discontinued immediately if serotonin syndrome symptoms occur and supportive symptomatic treatment should be initiated. If concomitant treatment with TONMYA and other serotonergic drugs is clinically warranted, careful observation is advised, particularly during treatment initiation or dosage increases.

Tricyclic antidepressant-like adverse reactions: Cyclobenzaprine is structurally related to TCAs. TCAs have been reported to produce arrhythmias, sinus tachycardia, prolongation of the conduction time leading to myocardial infarction and stroke. If clinically significant central nervous system (CNS) symptoms develop, consider discontinuation of TONMYA. Caution should be used when TCAs are given to patients with a history of seizure disorder, because TCAs may lower the seizure threshold. Patients with a history of seizures should be monitored during TCA use to identify recurrence of seizures or an increase in the frequency of seizures.

Atropine-like effects: Use with caution in patients with a history of urinary retention, angle-closure glaucoma, increased intraocular pressure, and in patients taking anticholinergic drugs.

CNS depression and risk of operating a motor vehicle or hazardous machinery: TONMYA monotherapy may cause CNS depression. Concomitant use of TONMYA with alcohol, barbiturates, or other CNS depressants may increase the risk of CNS depression. Advise patients not to operate a motor vehicle or dangerous machinery until they are reasonably certain that TONMYA therapy will not adversely affect their ability to engage in such activities. Oral mucosal adverse reactions: In clinical studies with TONMYA, oral mucosal adverse reactions occurred more frequently in patients treated with TONMYA compared to placebo. Advise patients to moisten the mouth with sips of water before administration of TONMYA to reduce the risk of oral sensory changes (hypoesthesia). Consider discontinuation of TONMYA if severe reactions occur.

ADVERSE REACTIONS

The most common adverse reactions (incidence $\geq 2\%$ and at a higher incidence in TONMYA-treated patients compared to placebo-treated patients) were oral hypoesthesia, oral discomfort, abnormal product taste, somnolence, oral paresthesia, oral pain, fatigue, dry mouth, and aphthous ulcer.

DRUG INTERACTIONS

MAO inhibitors: Life-threatening interactions may occur. Other serotonergic drugs: Serotonin syndrome has been reported. CNS depressants: CNS depressant effects of alcohol, barbiturates, and other CNS depressants may be enhanced. Tramadol: Seizure risk may be enhanced. Guanethidine or other similar acting drugs: The antihypertensive action of these drugs may be blocked.

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, TONMYA may cause fetal harm when administered to a pregnant woman. The limited amount of available observational data on oral cyclobenzaprine use in pregnancy is of insufficient quality to inform a TONMYA-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Advise pregnant women about the potential risk to the fetus with maternal exposure to TONMYA and to avoid use of TONMYA two weeks prior to conception and through the first trimester of pregnancy. Report pregnancies to the Tonix Medicines, Inc., adverse-event reporting line at 1-888-869-7633 (1-888-TNXPMED). Lactation: A small number of published cases report the transfer of cyclobenzaprine into human milk in low amounts, but these data cannot be confirmed. There are no data on the effects of cyclobenzaprine on a breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for TONMYA and any potential adverse effects on the breastfed child from TONMYA or from the underlying maternal condition. Pediatric use: The safety and effectiveness of TONMYA have not been established. Geriatric patients: Of the total number of TONMYA-treated patients in the clinical trials in adult patients with fibromyalgia, none were 65 years of age and older. Clinical trials of TONMYA did not include sufficient numbers of patients 65 years of age and older to determine whether they respond differently from younger adult patients. Hepatic impairment:

The recommended dosage of TONMYA in patients with mild hepatic impairment (HI) (Child Pugh A) is 2.8 mg once daily at bedtime, lower than the recommended dosage in patients with normal hepatic function. The use of TONMYA is not recommended in patients with moderate HI (Child Pugh B) or severe HI (Child Pugh C). Cyclobenzaprine exposure (AUC) was increased in patients with mild HI and moderate HI compared to subjects with normal hepatic function, which may increase the risk of TONMYA-associated adverse reactions.

Please see additional safety information in the full Prescribing Information.

To report suspected adverse reactions, contact Tonix Medicines, Inc. at 1-888-869-7633, or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.



Source: Tonix Pharmaceuticals Holding Corp.