December 22, 2022

Rigel Announces Availability of REZLIDHIA™ (olutasidenib) in the U.S. for the Treatment of Adult Patients with Relapsed or Refractory Acute Myeloid Leukemia with a Susceptible IDH1 Mutation

SOUTH SAN FRANCISCO, Calif., Dec. 22, 2022 /PRNewswire/ -- Rigel Pharmaceuticals, Inc. (Nasdaq: RIGL), today announced that REZLIDHIA™ (olutasidenib) capsules are available in the U.S. by prescription for the treatment of adult patients with relapsed or refractory (R/R) acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test. REZLIDHIA is an oral, small molecule, inhibitor of mutated IDH1 designed to bind to and inhibit mIDH1 to reduce 2-hydroxyglutarate levels and restore normal cellular differentiation of myeloid cells. REZLIDHIA was approved by the U.S. Food and Drug Administration (FDA) on December 1, 2022.

"We are pleased to launch REZLIDHIA today in the U.S. and are ready to provide access to patients," said Raul Rodriguez, Rigel's president and CEO. "REZLIDHIA is Rigel's second commercial product and is a new and promising oral treatment option for mIDH1 R/R AML patients who typically experience poor clinical outcomes. With durable efficacy data, promising overall survival, and a well characterized safety profile, we are excited to bring REZLIDHIA to patients in need."

The recommended dosage of REZLIDHIA is 150 mg taken orally twice daily. Based on the recommended dosage, the wholesale acquisition cost (WAC) of REZLIDHIA in the U.S. is $32,200 per month. REZLIDHIA is supplied as follows:

NDC Number	Product	Package	WAC
71332-005-01	Oral capsule 150 mg	Bottle - 30 capsules each	$16,100

To assist with access to REZLIDHIA, RIGEL ONECARE®, a comprehensive patient support center, can help patients and physicians as they navigate through insurance coverage

requirements and provide financial assistance when needed and if eligible, along with other support programs. Patients will be assigned a Nurse Navigator to assess the patient's individual case.

To learn more, visit www.REZLIDHIA.com or contact RIGEL ONECARE at 833-RIGELOC (833-744-3562).

In August 2022, Rigel and Forma Therapeutics, Inc. announced they entered an exclusive, worldwide license agreement to develop, manufacture and commercialize REZLIDHIA. Under the terms of the agreement, Rigel is responsible for the launch and commercialization of REZLIDHIA in the U.S., and intends to work with potential partners to further develop and commercialize the product outside the U.S.

About AML

Acute myeloid leukemia (AML) is a rapidly progressing cancer of the blood and bone marrow that affects myeloid cells, which normally develop into various types of mature blood cells. AML occurs primarily in adults and accounts for about 1 percent of all adult cancers. The American Cancer Society estimates that in the United States alone, there will be about 20,050 new cases, most in adults, in 2022.[1]

Relapsed AML affects about half of all patients who, following treatment and remission, experience a return of leukemia cells in the bone marrow.[2] Refractory AML, which affects between 10 and 40 percent of newly diagnosed patients, occurs when a patient fails to achieve remission even after intensive treatment.[3] Quality of life declines for patients with each successive line of treatment for AML, and well-tolerated treatments in relapsed or refractory disease remain an unmet need.

About REZLIDHIA™

INDICATION

REZLIDHIA is indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation as detected by an FDA-approved test.

IMPORTANT SAFETY INFORMATION

WARNING: DIFFERENTIATION SYNDROME

Differentiation syndrome, which can be fatal, can occur with REZLIDHIA treatment. Symptoms may include dyspnea, pulmonary infiltrates/pleuropericardial effusion, kidney injury, hypotension, fever, and weight gain. If differentiation syndrome is suspected, withhold REZLIDHIA and initiate treatment with corticosteroids and hemodynamic monitoring until symptom resolution.

WARNINGS AND PRECAUTIONS

Differentiation Syndrome

REZLIDHIA can cause differentiation syndrome. In the clinical trial of REZLIDHIA in patients with relapsed or refractory AML, differentiation syndrome occurred in 16% of patients, with grade 3 or 4 differentiation syndrome occurring in 8% of patients treated, and fatalities in 1% of patients. Differentiation syndrome is associated with rapid proliferation and differentiation of myeloid cells and may be life-threatening or fatal. Symptoms of differentiation syndrome in patients treated with REZLIDHIA included leukocytosis, dyspnea, pulmonary infiltrates/pleuropericardial effusion, kidney injury, fever, edema, pyrexia, and weight gain. Of the 25 patients who experienced differentiation syndrome, 19 (76%) recovered after

treatment or after dose interruption of REZLIDHIA. Differentiation syndrome occurred as early as 1 day and up to 18 months after REZLIDHIA initiation and has been observed with or without concomitant leukocytosis.

If differentiation syndrome is suspected, temporarily withhold REZLIDHIA and initiate systemic corticosteroids (e.g., dexamethasone 10 mg IV every 12 hours) for a minimum of 3 days and until resolution of signs and symptoms. If concomitant leukocytosis is observed, initiate treatment with hydroxyurea, as clinically indicated. Taper corticosteroids and hydroxyurea after resolution of symptoms. Differentiation syndrome may recur with premature discontinuation of corticosteroids and/or hydroxyurea treatment. Institute supportive measures and hemodynamic monitoring until improvement; withhold dose of REZLIDHIA and consider dose reduction based on recurrence.

Hepatotoxicity
REZLIDHIA can cause hepatotoxicity, presenting as increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), increased blood alkaline phosphatase, and/or elevated bilirubin. Of 153 patients with relapsed or refractory AML who received REZLIDHIA, hepatotoxicity occurred in 23% of patients; 13% experienced grade 3 or 4 hepatotoxicity. One patient treated with REZLIDHIA in combination with azacitidine in the clinical trial, a combination for which REZLIDHIA is not indicated, died from complications of drug-induced liver injury. The median time to onset of hepatotoxicity in patients with relapsed or refractory AML treated with REZLIDHIA was 1.2 months (range: 1 day to 17.5 months) after REZLIDHIA initiation, and the median time to resolution was 12 days (range: 1 day to 17 months). The most common hepatotoxicities were elevations of ALT, AST, blood alkaline phosphatase, and blood bilirubin.

Monitor patients frequently for clinical symptoms of hepatic dysfunction such as fatigue, anorexia, right upper abdominal discomfort, dark urine, or jaundice. Obtain baseline liver function tests prior to initiation of REZLIDHIA, at least once weekly for the first two months, once every other week for the third month, once in the fourth month, and once every other month for the duration of therapy. If hepatic dysfunction occurs, withhold, reduce, or permanently discontinue REZLIDHIA based on recurrence/severity.

ADVERSE REACTIONS
The most common (≥20%) adverse reactions, including laboratory abnormalities, were aspartate aminotransferase increased, alanine aminotransferase increased, potassium decreased, sodium decreased, alkaline phosphatase increased, nausea, creatinine increased, fatigue/malaise, arthralgia, constipation, lymphocytes increased, bilirubin increased, leukocytosis, uric acid increased, dyspnea, pyrexia, rash, lipase increased, mucositis, diarrhea and transaminitis.

DRUG INTERACTIONS

- Avoid concomitant use of REZLIDHIA with strong or moderate CYP3A inducers.
- Avoid concomitant use of REZLIDHIA with sensitive CYP3A substrates unless otherwise instructed in the substrates prescribing information. If concomitant use is unavoidable, monitor patients for loss of therapeutic effect of these drugs.

LACTATION
Advise women not to breastfeed during treatment with REZLIDHIA and for 2 weeks after the

last dose.

GERIATRIC USE
No overall differences in effectiveness were observed between patients 65 years and older and younger patients. Compared to patients younger than 65 years of age, an increase in incidence of hepatotoxicity and hypertension was observed in patients ≥65 years of age.

HEPATIC IMPAIRMENT
In patients with mild or moderate hepatic impairment, closely monitor for increased probability of differentiation syndrome.

Click here for Full Prescribing Information, including Boxed WARNING.

To report side effects of prescription drugs to the FDA, visitwww.fda.gov/medwatch or call 1-800-FDA-1088 (800-332-1088).

REZLIDHIA is a trademark and RIGEL ONECARE is a registered trademark of Rigel Pharmaceuticals, Inc.

RIGEL ONECARE is a patient support center sponsored by Rigel Pharmaceuticals, Inc.

About Rigel
Rigel Pharmaceuticals, Inc. (Nasdaq: RIGL) is a biotechnology company dedicated to discovering, developing and providing novel small molecule drugs that significantly improve the lives of patients with hematologic disorders, cancer, and rare immune diseases. Founded in 1996, Rigel is based in South San Francisco, California. For more information on Rigel, the Company's marketed products and pipeline of potential products, visit www.rigel.com.

1. The American Cancer Society. Key Statistics for Acute Myeloid Leukemia (AML). Revised January 12, 2022. Accessed Aug. 1, 2022 at https://www.cancer.org/cancer/acute-myeloid-leukemia/about/key-statistics.html
2. Leukaemia Care. (2019). Relapse in Acute Myeloid Leukaemia (AML). Version 3. Reviewed October 2021. Accessed Dec 2, 2021 at https://media.leukaemiacare.org.uk/wp-content/uploads/Relapse-in-Acute-Myeloid-Leukaemia-AML-Web-Version.pdf
3. Thol F, Schlenk RF, Heuser M, Ganser A. How I treat refractory and early relapsed acute myeloid leukemia. Blood. 2015 Jul 16;126(3):319-27. doi: https://doi.org/10.1182/blood-2014-10-551911
4. REZLIDHIA™ [package insert] South San Francisco, CA: Rigel Pharmaceuticals, Inc.

Forward Looking Statements
This press release contains forward-looking statements relating to, among other things, our ability to fill prescriptions of REZLIDHIA, the availability and cost of REZLIDHIA, and the benefit of REZLIDHIA to people who typically experience poor clinical outcomes. Any statements contained in this press release that are not statements of historical fact may be deemed to be forward-looking statements. Forward-looking statements can be identified by words such as "plan", "potential", "may", "expects", "will" and similar expressions in reference to future periods. Forward-looking statements are neither historical facts nor assurances of future performance. Instead, they are based on Rigel's current beliefs,

expectations, and assumptions regarding the future of our business, future plans and strategies, projections, anticipated events and trends, the economy and other future conditions, and hence they inherently involve significant risks, uncertainties and changes in circumstances that are difficult to predict and many of which are outside of our control. Therefore, you should not rely on any of these forward-looking statements. Actual results and the timing of events could differ materially from those anticipated in such forward-looking statements as a result of these risks and uncertainties, which include, without limitation, risks that the FDA, European Medicines Agency or other regulatory authorities may make adverse decisions regarding olutasidenib; risks that clinical trials may not be predictive of real-world results or of results in subsequent clinical trials; risks that olutasidenib may have unintended side effects, adverse reactions or incidents of misuses; the availability of resources to develop Rigel's product candidates; market competition; as well as other risks detailed from time to time in Rigel's reports filed with the Securities and Exchange Commission, including its Quarterly Report on Form 10-Q for the quarter ended September 30, 2022 and subsequent filings. Any forward-looking statement made by us in this press release is based only on information currently available to us and speaks only as of the date on which it is made. Rigel does not undertake any obligation to update forward-looking statements, whether written or oral, that may be made from time to time, whether as a result of new information, future developments or otherwise, and expressly disclaims any obligation or undertaking to release publicly any updates or revisions to any forward-looking statements contained herein, except as required by law.

Rigel Contacts
Media:
David Rosen, Argot Partners
Phone: 212.600.1902
Email: david.rosen@argotpartners.com

Investors:
Rigel Pharmaceuticals, Inc.
Phone: 650.624.1232
Email: ir@rigel.com

View original content to download multimedia:https://www.prnewswire.com/news-releases/rigel-announces-availability-of-rezlidhia-olutasidenib-in-the-us-for-the-treatment-of-adult-patients-with-relapsed-or-refractory-acute-myeloid-leukemia-with-a-susceptible-idh1-mutation-301708714.html

SOURCE Rigel Pharmaceuticals, Inc.