



Phase 2 Clinical Trial To Evaluate The Efficacy Of Phentolamine Ophthalmic Solution And Low-Dose Pilocarpine For The Treatment Of Presbyopia

(Paper ID 76645)

Presenter: Jay S. Pepose, MD, PhD, ABO

Authors:

Jay S. Pepose¹, MD, PhD, ABO; Mina P. Sooch, BEng, MBA²; Konstantinos Charizanis, PhD, MBA⁴; Charles B. Slonim, MD³; Alan R. Meyer, MBA⁴; Eliot Lazar, MD¹

Author affiliations:

1. Medical Advisor to Ocuphire Pharma, Inc
2. Employed by Ocuphire Pharma, Inc
3. Employed by Oculos Development Services, LLC
4. Consultant to Ocuphire Pharma, Inc

Dr. Pepose's financial disclosures:

- Acufocus
- Allergan
- Bausch Health (Valeant)
- GlaxoSmithKline
- Johnson and Johnson Vision
- Keeler
- Novartis
- Ocunexus
- Ocuphire
- Okogen
- Stuart Pharmaceuticals
- Sun Pharma
- TearLab
- Thea Pharm


Phentolamine Ophth. Solution (POS) + Low-Dose Pilocarpine (LDP) Combination

Moderate Use Of Iris Dilator And Iris Sphincter Muscles To Improve Near Vision

0.75% POS



**Iris Dilator
Muscle
Inhibition**




**Iris Sphincter
Muscle
Activation**



0.4% LDP

- Alpha1/2 antagonist approved decades ago 505(b)(2)
- Novel MOA on iris dilator with 24+ hour durability with moderate 1+mm pupil reduction
- No daytime redness with chronic evening dosing of POS
- Well-tolerated with no systemic effects
- Stable, preservative-free, single use vial




*1.5 to 2.5 mm PD reduction
moves toward the pin-hole
(1.6 to 2.5 mm, up to <3 mm)*

- Cholinergic agonist approved decades ago 505(b)(2)
- Known MOA on sphincter muscle with more potent miotic effects at approved doses (1%, 2%, 4%)
- Chronic daily dosing of LDP in daytime
- Low concentration avoids known tolerability issues:
 - headache and browache
 - redness
 - accommodative spasm causing loss of distance vision especially at night

Presbyopia VEGA-1 Phase 2 Design

Randomized, Double-Masked, Placebo-Controlled Multi-Center One-Week Trial



Eligibility Criteria

- Males or females ≥ 40 and ≤ 64 years of age
- BCDVA of 0.0 LogMAR (20/20 Snellen equivalent) or better in each eye under photopic conditions
- DCNVA of 0.4 LogMAR (20/50 Snellen equivalent) or worse under photopic conditions in each eye & binocularly

Endpoints

Primary: % of subjects with ≥ 3 lines of improvement in distance-corrected near visual acuity comparing POS + LDP vs placebo alone at 1 hour

Secondary:

- % of subjects with ≥ 2 and ≥ 3 lines gained at time points from 30 min to 6 hours in photopic and mesopic lighting comparing POS + LDP vs placebo, POS alone, and LDP alone
- No loss of distance vision
- Pupil diameter at time points
- Safety and tolerability (redness)

Demographics and Baseline Characteristics



Treatment And Placebo Arms Were Balanced In the VEGA-1 Phase 2 Clinical Trial

	Placebo Alone N=43	POS Alone N=30	LDP Alone N=31	POS+LDP N=43	Total N=147
Age (years): Median (Range)	52 (42-62)	54 (41-60)	52 (44-64)	53 (43-63)	53 (41-64)
Sex: Male n (%)	15 (35%)	7 (23%)	13 (42%)	5 (12%)	40 (27%)
Female n (%)	28 (65%)	23 (77%)	18 (58%)	38 (88%)	107 (73%)
Race: White n (%)	37 (86%)	26 (87%)	28 (90%)	40 (93%)	131 (89%)
Other* n (%)	6 (14%)	1 (3%)	3 (10%)	3 (7%)	15 (11%)
Dark Iris Color: n (%)	18 (42%)	12 (40%)	12 (39%)	18 (42%)	60 (41%)
Light Iris Color: n (%)	25 (58%)	18 (60%)	19 (61%)	25.1 (58%)	87 (59%)
Photopic DCNVA Mean Letters read- Binocular (Snellen Equiv.) <i>70 letters = 20/20</i>	46 (20/63)	45 (20/63)	48 (20/63)	46 (20/63)	46 (20/63)
Photopic BCDVA Mean Letters read- Binocular (Snellen Equiv.) <i>55 letters = 20/20</i>	62 (20/15)	61 (20/15)	60 (20/15)	61 (20/15)	61 (20/15)
Photopic Pupil Diameter Mean (mm)	4.3	4.5	4.3	4.3	4.3
Mesopic Pupil Diameter Mean (mm)	5.1	5.0	5.0	5.1	5.1
IOP (mmHg)	13.5	14.8	13.9	14.4	14.1

Primary Endpoint: % Of Subjects ≥ 15 Letter Gain In Photopic DCNVA At 1 Hour

Primary Endpoint Was Met For POS + LDP Gaining ≥ 15 Letters Near Vision In PP Population

VEGA-1 Phase 2 Trial




Note: PP population differs from mITT by only one subject; results were essentially identical.

Source: VEGA-1 TLR Table 14.2.1.2 %of Subjects With Improvement From Baseline in Photopic DCNVA by Time Point (PP Population). 15 letters is 3 lines and 10 letters is 2 lines.

Secondary Endpoint: % Of Subjects ≥ 15 Letter Gain At All Timepoints

POS + LDP Had Strong Response With ≥ 15 Letter Gain From 30 Min To 6 Hours





Source: VEGA-1 TLR Table 14.2.1.2 Percent of Subjects with Improvement From Baseline in Photopic DCNVA by Time Point (PP Population). 15 letters is 3 lines.

Secondary Endpoint: Snellen Equiv. DCNVA Baseline vs. 1 Hour Post POS+LDP

POS+LDP Had A Dramatic Effect On Near Vision With A Clear Improvement Shift

VEGA-1 Phase 2 Trial

Binocular DCNVA
POS + LDP




26% Placebo Adjusted Response $\leq 20/40$ for POS+LDP Combo at 1 Hour ($p=0.02$) Compared to AGN
13% Placebo Adjusted ($p= 0.02$) photopic DCNVA

Source: VEGA-1 TLR Table 14.2.24.1 Percent of Subjects with Photopic DCNVA by Time Point (PP Population); Novartis/Encore Materials, ASCRS 2021 Allergan Abstracts



Secondary Endpoint: % of Subjects ≥ 15 Letter Gain In Near & ≤ 5 Letter Loss In Distance

Phase 3 Approval Endpoint Also Showed Early Onset & Durable Near Vision Gain Without Loss Of Distance




Secondary Endpoint: Change In Photopic/Mesopic BCDVA At 1 Hour


Treatment With POS And/Or LDP Did Not Reduce BCDVA And Had A Modest Beneficial Effect

VEGA-1 Phase 2 Trial

Percent of Subjects With Improvement or Loss From Baseline in Photopic BCDVA at 1 hour




Percent of Subjects With Improvement or Loss From Baseline in Mesopic BCDVA at 1 hour



Secondary Endpoint: Mean Pupil Diameter Over Time

Achieved Pupil Size ~2mm In POS+LDP Consistent With 3-line Improvement In Near Vision



Source: VEGA-1 TLR Table 14.2.12.1 Observed Values and Change from Baseline in Photopic Pupil Diameter by Time Point (PP Population)



Secondary Endpoint: Safety Findings

POS+LDP Combination Was Well Tolerated With A Favorable Safety Profile In VEGA-1 Trial

	Placebo Alone n=45	POS Alone n=30	LDP Alone n=31	POS+LDP n=44
Total Treatment Emergent Adverse Events (n)	4	18	13	50
TEAEs by Severity (n [%])				
Mild	1 (2.2%)	6 (20%)	6 (19.4%)	13 (29.5%)
Moderate	1 (2.2%)	0 (0%)	0 (0%)	1 (2.3%)
Severe	0 (0%)	0 (0%)	0 (0%)	1 (2.3%)
AEs Occurring in ≥ 5% of subjects (n [%])				
Instillation Site Pain (Mild)	1 (2.2%)	3 (10%)	0 (0%)	4 (9.1%)
Instillation Site Erythema (Mild)	0 (0%)	3 (10%)	2 (6.5%)	5 (11.4%)
Conjunctival Hyperemia (Mild)	0 (0%)	2 (6.7%)	0 (0%)	2 (4.5%)
Eye Disorders (Mild)	1 (2.2%)	2 (6.7%)	4 (12.9%)	5 (11.4%)

- No deaths, no serious AEs, and 1 withdrawal due to AEs (on POS alone)
- **0% Headaches or Browaches reported for POS+LDP and POS alone**
- Only 1 subject in LDP alone arm reported mild headache
- Almost all AEs were mild and most common was mild instillation site discomfort
- **Distance visual acuity not adversely affected (as shown earlier)**
- No change in IOP

Conjunctival Hyperemia CCLRU Scale for Reference



Summary Of Positive VEGA-1 Phase 2 Results

POS + LDP Efficacy Data With A Favorable Safety Profile In Presbyopia

- **Met the primary endpoint** with statistical significance for binocular photopic near vision at 1 hour
 - 61% POS+ LDP gained 15 letters (3 lines) or more vs. 28% Placebo (33% Placebo Adjusted)
- **Met the Phase 3 co-primary endpoint** vs. placebo gaining 15 letters (3 lines) near vision with less than 5 letters of distance vision loss
- **Met many key secondary endpoints**
 - Rapid onset at 30 min
 - Durable near vision improvement through at least 6 hours
 - POS+LDP was numerically better than each component through 2-hours
 - A majority of subjects treated with the combination achieved near acuity of 20/30 or better
 - Sustained significant reduction in PD over at least 18 hours due the durability effects of POS
 - Near vision efficacy seen monocularly and binocularly
 - Also, efficacy data in both light and dark iris colors
- **Favorable safety profile for POS + LDP**
 - No serious AEs
 - No systemic AEs were observed in >5% subjects
 - No headaches, no browaches, and no blurry vision AEs were reported
 - Only mild, transient conjunctival hyperemia observed in <5% of subjects
- **Positive Phase 2 results lead to advancing Phase 3 presbyopia program**

Thank you for your attention!

DR. JAY S. PEPOSE

Medical Director
Pepose Vision Institute
Professor of Clinical Ophthalmology And Visual Sciences
Washington University School of Medicine In St. Louis
St. Louis, MO, United States
Email: jpepose@peposevision.com

VEGA-1 Clinical Trial Sponsor is Ocuphire Pharma
www.Ocuphire.com