

September 18, 2025



# **Tonix Pharmaceuticals Announces Positive Pre-IND Meeting with FDA for TNX-102 SL for the Treatment of Major Depressive Disorder**

*Tonix anticipates filing the IND application in the fourth quarter of 2025*

*TNX-102 SL is a potential first-in-class treatment for targeting the disturbed sleep associated with depression*

*TNX-102 SL is FDA-approved for the treatment of fibromyalgia*

CHATHAM, N.J., Sept. 18, 2025 (GLOBE NEWSWIRE) -- Tonix Pharmaceuticals Holding Corp. (Nasdaq: TNXP), a fully-integrated biotechnology company with marketed products and a pipeline of development candidates, today announced the successful completion of a Type B Pre-Investigational New Drug (Pre-IND) meeting with the U.S. Food and Drug Administration (FDA) regarding the development of TNX-102 SL (sublingual cyclobenzaprine HCl) for the treatment of major depressive disorder (MDD). The Company received positive feedback from the FDA and plans to pursue a supplemental new drug application (sNDA) to expand the therapeutic indication of TNX-102 SL to include MDD, based on exploratory findings suggesting that improving sleep quality may positively impact depressive symptoms.

"We are pleased with the outcome of our Pre-IND meeting with the FDA and appreciate their thoughtful guidance," said Seth Lederman, M.D., Chief Executive Officer of Tonix Pharmaceuticals. "This marks a significant step forward in our efforts to develop TNX-102 SL as a novel treatment for MDD, a condition that affects millions and remains underserved by current therapies that are often difficult to tolerate."

"We believe bedtime TNX-102 SL has the potential to be a first-in-class treatment that is designed to target the reduced quality and quantity of slow wave sleep associated with depression," said Dr. Gregory Sullivan, M.D., Chief Medical Officer of Tonix Pharmaceuticals. "TNX-102 SL is a tertiary amine tricyclic that is designed for transmucosal absorption to bypass first pass hepatic metabolism. In contrast, FDA-approved tertiary amine tricyclic antidepressants are swallowed pill formulations, that are largely metabolized by first-pass to longer-lived secondary amine tricyclics. Also, the FDA-approved tertiary amine tricyclic antidepressants are only active in treating MDD at more than ten times the dose employed with TNX-102 SL and can adversely impact weight, blood pressure, cognition, and sexual function."

The FDA provided constructive feedback during the Pre-IND meeting for TNX-102 SL in

MDD and found the proposed long-term safety data collection plan generally reasonable, potentially streamlining the development path.

An IND filing is planned for Q4 2025, positioning the program to enter Phase 2 clinical trials shortly thereafter. TNX-102 SL is FDA approved for the treatment of fibromyalgia, under the brand name Tonmya™. In the Phase 3 RESILIENT study of fibromyalgia patients, the TNX-102 SL – treated group had activity on improving depression over placebo by the Beck Depression Inventory (BDI), with an uncorrected p-value < 0.05. The biological relationship between depressed symptoms in fibromyalgia and those in MDD is not understood.

### **About TNX-102 SL**

TNX-102 SL is a patented sublingual tablet formulation of cyclobenzaprine hydrochloride that enables rapid transmucosal absorption and reduces production of the long half-life active metabolite, norcyclobenzaprine, by bypassing first-pass hepatic metabolism. As a tertiary amine tricyclic (TAT) and multifunctional agent with potent binding and antagonist activities at the 5-HT<sub>2A</sub> serotonergic, α<sub>1</sub>-adrenergic, H<sub>1</sub>-histaminergic, and M<sub>1</sub>-muscarinic receptors. It is currently approved in the U.S. as a once-daily bedtime treatment for fibromyalgia in adults. TNX-102 SL is also in development as a daily bedtime treatment for acute stress reaction/acute stress disorder. Tonix also holds active IND's for the following indications for TNX-102 SL: Long COVID (post-acute sequelae of COVID-19), PTSD, alcohol use disorder, and agitation in Alzheimer's disease. The United States Patent and Trademark Office (USPTO) issued United States Patent No. 9636408 in May 2017, Patent No. 9956188 in May 2018, Patent No. 10117936 in November 2018, Patent No. 10357465 in July 2019, and Patent No. 10736859 in August 2020. The Protectic™ protective eutectic and Angstro-Technology™ formulation claimed in the patent are important elements of Tonix's proprietary composition. These patents are expected to provide Tonmya with U.S. market exclusivity until 2034. Pending patent applications related to method of use could extend exclusivity until 2044.

### **Tonix Pharmaceuticals Holding Corp.**

Tonix Pharmaceuticals is a fully-integrated biotechnology company with marketed products and a pipeline of development candidates. Tonix recently received FDA approval for Tonmya™, a first-in-class, non-opioid analgesic medicine for the treatment of fibromyalgia, a chronic pain condition that affects millions of adults. This marks the first approval for a new prescription medicine for fibromyalgia in more than 15 years. Tonix also markets two treatments for acute migraine in adults. Tonix's development portfolio is focused on central nervous system (CNS) disorders, immunology, immuno-oncology and infectious diseases. TNX-102 SL is being developed to treat acute stress reaction and acute stress disorder under a Physician-Initiated IND at the University of North Carolina in the OASIS study funded by the U.S. Department of Defense (DoD). Tonix's immunology development portfolio consists of biologics to address organ transplant rejection, autoimmunity and cancer, including TNX-1500, which is an Fc-modified humanized monoclonal antibody targeting CD40-ligand (CD40L or CD154) being developed for the prevention of allograft rejection and for the treatment of autoimmune diseases. Tonix's infectious disease portfolio includes TNX-801, a vaccine in development for mpox and smallpox, as well as TNX-4200 for which Tonix has a contract with the U.S. DoD's Defense Threat Reduction Agency (DTRA) for up to \$34 million over five years. TNX-4200 is a small molecule broad-spectrum antiviral agent targeting CD45 for the prevention or treatment of infections to improve the

medical readiness of military personnel in biological threat environments. Tonix owns and operates a state-of-the art infectious disease research facility in Frederick, Md.

\* Tonix's product development candidates are investigational new drugs or biologics; their efficacy and safety have not been established and have not been approved for any indication.

This press release and further information about Tonix can be found at [www.tonixpharma.com](http://www.tonixpharma.com).

### **Forward Looking Statements**

Certain statements in this press release are forward-looking within the meaning of the Private Securities Litigation Reform Act of 1995. These statements may be identified by the use of forward-looking words such as "anticipate," "believe," "forecast," "estimate," "expect," and "intend," among others. These forward-looking statements are based on Tonix's current expectations and actual results could differ materially. There are a number of factors that could cause actual events to differ materially from those indicated by such forward-looking statements. These factors include, but are not limited to, risks related to the failure to successfully launch and commercialize Tonmya and any of our approved products; risks related to the failure to obtain FDA clearances or approvals and noncompliance with FDA regulations; risks related to the failure to successfully launch and commercialize Tonmya and any of our approved products; risks related to the timing and progress of clinical development of our product candidates; our need for additional financing; uncertainties of patent protection and litigation; uncertainties of government or third party payor reimbursement; limited research and development efforts and dependence upon third parties; and substantial competition. As with any pharmaceutical under development, there are significant risks in the development, regulatory approval and commercialization of new products. Tonix does not undertake an obligation to update or revise any forward-looking statement. Investors should read the risk factors set forth in the Annual Report on Form 10-K for the year ended December 31, 2024, as filed with the Securities and Exchange Commission (the "SEC") on March 18, 2025, and periodic reports filed with the SEC on or after the date thereof. All of Tonix's forward-looking statements are expressly qualified by all such risk factors and other cautionary statements. The information set forth herein speaks only as of the date thereof.

### **Investor Contact**

Jessica Morris  
Tonix Pharmaceuticals  
[investor.relations@tonixpharma.com](mailto:investor.relations@tonixpharma.com)  
(862) 799-8599

Brian Korb  
astr partners  
[brian.korb@astrpartners.com](mailto:brian.korb@astrpartners.com)  
(917) 653-5122

### **Media Contact**

Ray Jordan  
Putnam Insights  
[ray@putnaminsights.com](mailto:ray@putnaminsights.com)

(949) 245-5432

## INDICATION

TONMYA is indicated for the treatment of fibromyalgia in adults.

## CONTRAINDICATIONS

TONMYA is contraindicated:

In patients with hypersensitivity to cyclobenzaprine or any inactive ingredient in TONMYA. Hypersensitivity reactions may manifest as an anaphylactic reaction, urticaria, facial and/or tongue swelling, or pruritus. Discontinue TONMYA if a hypersensitivity reaction is suspected.

With concomitant use of monoamine oxidase (MAO) inhibitors or within 14 days after discontinuation of an MAO inhibitor. Hyperpyretic crisis seizures and deaths have occurred in patients who received cyclobenzaprine (or structurally similar tricyclic antidepressants) concomitantly with MAO inhibitors drugs.

During the acute recovery phase of myocardial infarction, and in patients with arrhythmias, heart block or conduction disturbances, or congestive heart failure.

In patients with hyperthyroidism.

## WARNINGS AND PRECAUTIONS

**Embryofetal toxicity:** Based on animal data, TONMYA may cause neural tube defects when used two weeks prior to conception and during the first trimester of pregnancy. Advise females of reproductive potential of the potential risk and to use effective contraception during treatment and for two weeks after the final dose. Perform a pregnancy test prior to initiation of treatment with TONMYA to exclude use of TONMYA during the first trimester of pregnancy.

**Serotonin syndrome:** Concomitant use of TONMYA with selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, tramadol, bupropion, meperidine, verapamil, or MAO inhibitors increases the risk of serotonin syndrome, a potentially life-threatening condition. Serotonin syndrome symptoms may include mental status changes, autonomic instability, neuromuscular abnormalities, and/or gastrointestinal symptoms. Treatment with TONMYA and any concomitant serotonergic agent should be discontinued immediately if serotonin syndrome symptoms occur and supportive **symptomatic treatment should be initiated**. If concomitant treatment with TONMYA and other serotonergic drugs is clinically warranted, careful observation is advised, particularly during treatment initiation or dosage increases.

**Tricyclic antidepressant-like adverse reactions:** Cyclobenzaprine is structurally related to TCAs. TCAs have been reported to produce arrhythmias, sinus tachycardia, prolongation of the conduction time leading to myocardial infarction and stroke. If clinically significant central nervous system (CNS) symptoms develop, consider discontinuation of TONMYA. Caution should be used when TCAs are given to patients with a history of seizure disorder, because TCAs may lower the seizure threshold. Patients with a history of seizures should be monitored during TCA use to identify recurrence of seizures or an increase in the frequency of seizures.

**Atropine-like effects:** Use with caution in patients with a history of urinary retention, angle-closure glaucoma, increased intraocular pressure, and in patients taking anticholinergic drugs.

CNS depression and risk of operating a motor vehicle or hazardous machinery: TONMYA monotherapy may cause CNS depression. Concomitant use of TONMYA with alcohol, barbiturates, or other CNS depressants may increase the risk of CNS depression. Advise patients not to operate a motor vehicle or dangerous machinery until they are reasonably certain that TONMYA therapy will not adversely affect their ability to engage in such activities.

Oral mucosal adverse reactions: In clinical studies with TONMYA, oral mucosal adverse reactions occurred more frequently in patients treated with TONMYA compared to placebo. Advise patients to moisten the mouth with sips of water before administration of TONMYA to reduce the risk of oral sensory changes (hypoesthesia). Consider discontinuation of TONMYA if severe reactions occur.

### **ADVERSE REACTIONS**

The most common adverse reactions (incidence  $\geq 2\%$  and at a higher incidence in TONMYA-treated patients compared to placebo-treated patients) were oral hypoesthesia, oral discomfort, abnormal product taste, somnolence, oral paresthesia, oral pain, fatigue, dry mouth, and aphthous ulcer.

### **DRUG INTERACTIONS**

MAO inhibitors: Life-threatening interactions may occur.

Other serotonergic drugs: Serotonin syndrome has been reported.

CNS depressants: CNS depressant effects of alcohol, barbiturates, and other CNS depressants may be enhanced.

Tramadol: Seizure risk may be enhanced.

Guanethidine or other similar acting drugs: The antihypertensive action of these drugs may be blocked.

### **USE IN SPECIFIC POPULATIONS**

Pregnancy: Based on animal data, TONMYA may cause fetal harm when administered to a pregnant woman. The limited amount of available observational data on oral cyclobenzaprine use in pregnancy is of insufficient quality to inform a TONMYA-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Advise pregnant women about the potential risk to the fetus with maternal exposure to TONMYA and to avoid use of TONMYA two weeks prior to conception and through the first trimester of pregnancy. Report pregnancies to the Tonix Medicines, Inc., adverse-event reporting line at 1-888-869-7633 (1-888-TNXPMED).

Lactation: A small number of published cases report the transfer of cyclobenzaprine into human milk in low amounts, but these data cannot be confirmed. There are no data on the effects of cyclobenzaprine on a breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for TONMYA and any potential adverse effects on the breastfed child from TONMYA or from the underlying maternal condition.

Pediatric use: The safety and effectiveness of TONMYA have not been established.

Geriatric patients: Of the total number of TONMYA-treated patients in the clinical trials in adult patients with fibromyalgia, none were 65 years of age and older. Clinical trials of TONMYA did not include sufficient numbers of patients 65 years of age and older to determine whether they respond differently from younger adult patients.

Hepatic impairment: The recommended dosage of TONMYA in patients with mild hepatic impairment (HI) (Child Pugh A) is 2.8 mg once daily at bedtime, lower than the recommended dosage in patients with normal hepatic function. The use of TONMYA is not recommended in patients with moderate HI (Child Pugh B) or severe HI (Child Pugh C). Cyclobenzaprine exposure (AUC) was increased in patients with mild HI and moderate HI compared to subjects with normal hepatic function, which may increase the risk of TONMYA-associated adverse reactions.

**Please see additional safety information in the full Prescribing Information.**

**To report suspected adverse reactions, contact Tonix Medicines, Inc. at 1-888-869-7633, or the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**



Source: Tonix Pharmaceuticals Holding Corp.